



Department of Economics and Management

**DEM Working Paper Series**

**Independent Factor Autoregressive Conditional Density Model**

Alexios Ghalanos  
(Cass Business School)

Eduardo Rossi  
(Università di Pavia)

Giovanni Urga  
(Cass Business School e Università di Bergamo)

**# 21 (11-12)**

Via San Felice, 5  
I-27100 Pavia  
<http://epmq.unipv.eu/site/home.html>

**November 2012**

# Independent Factor Autoregressive Conditional Density Model

Alexios Ghalanos <sup>\*</sup>                      Eduardo Rossi <sup>†</sup>  
Cass Business School, UK              University of Pavia, Italy

Giovanni Urga <sup>‡</sup>  
Cass Business School, UK and Bergamo University, Italy

November 12, 2012

## Abstract

In this paper, we propose a novel Independent Factor Autoregressive Conditional Density (IFACD) model able to generate time-varying higher moments using an independent factor setup. Our proposed framework incorporates dynamic estimation of higher comovements and feasible portfolio representation within a non elliptical multivariate distribution. We report an empirical application, using returns data from 14 MSCI equity index iShares for the period 1996 to 2011, and we show that the IFACD model provides superior VaR forecasts and portfolio allocations with respect to the CHICAGO and DCC models.

Keywords: Independent Factor Model, GO-GARCH, Independent Component Analysis, Time-varying Co-moments.

JEL Classification: C13, C16, C32, G11.

---

<sup>\*</sup>Faculty of Finance, 106 Bunhill Row, London EC1Y 8TZ, UK. [alexios.ghalanos.1@cass.city.ac.uk](mailto:alexios.ghalanos.1@cass.city.ac.uk).

<sup>†</sup>Dipartimento di Scienze Economiche e Aziendali. Via San Felice 5, 27100 Pavia, Italy. [erossi@eco.unipv.it](mailto:erossi@eco.unipv.it).

<sup>‡</sup>Faculty of Finance, 106 Bunhill Row, London EC1Y 8TZ, UK. [g.urga@city.ac.uk](mailto:g.urga@city.ac.uk).

# Independent Factor Autoregressive Conditional Density Model

## Abstract

In this paper, we propose a novel Independent Factor Autoregressive Conditional Density (IFACD) model able to generate time-varying higher moments using an independent factor setup. Our proposed framework incorporates dynamic estimation of higher comovements and feasible portfolio representation within a non elliptical multivariate distribution. We report an empirical application, using returns data from 14 MSCI equity index iShares for the period 1996 to 2011, and we show that the IFACD model provides superior VaR forecasts and portfolio allocations with respect to the CHICAGO and DCC models.

Keywords: Independent Factor Model, GO-GARCH, Independent Component Analysis, Time-varying Co-moments.

JEL Classification: C16, C32, G11.

## 1 INTRODUCTION

This paper develops an independent factor model for the conditional density of financial returns with time-varying skewness and kurtosis. In the presence of nonnormally distributed asset returns, optimal portfolio selection techniques require estimates of variance-covariance parameters, along with estimates of higher-order moments and co-moments of returns. The importance in portfolio analysis of modelling higher conditional moment dynamics has been highlighted by, *inter alia*, Barone Adesi et al. (2004), Beaulieu et al. (2007, 2009), Chabi-Yo et al. (2008), Harvey and Siddique (1999), Martellini and Ziemann (2010), Mencia and Sentana (2009, 2012), Sentana (2009), and Wilhelmsson (2009). In the univariate context, models that include higher moments dynamics have received growing attention. See for example Jondeau and Rockinger (2003, 2009), Harvey and Siddique (1999), Rockinger and Jondeau (2002), and Wilhelmsson (2009). However, little effort has been devoted to incorporate such dynamics in a multivariate context, mainly because of the difficulty to parameterize marginal and joint distributional parameters. There are very few notable exceptions. Mencia and Sentana (2009) adopt a flexible family of multivariate asymmetric distributions, known as location-scale mixtures of normals, which nests as particular cases several important elliptical symmetric distributions. When the distribution of asset returns can be expressed as a location-scale mixtures of normals, then the portfolio conditional distribution has time-varying higher moments arising from the interaction of the dynamics of the location, scale and skew parameters. On the other hand, Jondeau and Rockinger (2009) propose an asymmetric DCC-skew-T model with higher moment dynamics, but the presence of the skew and shape parameters in the conditional likelihood makes the estimation feasible only for a few assets. Broda and Paoletta (2009) propose a Conditionally Heteroskedastic Independent Component Analysis of Generalized Orthogonal GARCH of van der Weide (2002), the CHICAGO model, which is based on a multivariate affine representation of the Normal Inverse Gaussian (maNIG), originally proposed by Schmidt et al. (2006). In the CHICAGO model higher moments of the conditional distribution of asset returns are explicitly modelled albeit they are constant. The estimation of the CHICAGO model is based on the Independent Components Analysis (ICA) method used in Chen et al. (2008) and Zhang and Chan (2009). Unlike other models, independence offers a greater deal of flexibility in modelling the full marginal dynamics within a multivariate affine factor framework, enabling the calculation of conditional portfolio density used in risk management applications.

In this paper, we propose an Independent Factor Autoregressive Conditional Density model (IFACD)

which extends the CHICAGO model by allowing for a flexible parametrization of the dynamics of factor higher moments. The marginal density of each independent factor is supposed to be a one-dimensional Generalized Hyperbolic (GH) distribution, with skew and shape parameters modelled with autoregressive dynamics, as in the Autoregressive Conditional Density (ACD) model of Hansen (1994). This entails that the conditional density of returns is the multivariate affine GH (maGH) of Schmidt et al. (2006). An interesting implication of this factor representation is that conditional higher moments of asset returns and portfolios have a closed-form expression, and the portfolio conditional density can be obtained by Fast Fourier Transform (FFT).

The paper is organized as follows. Section 2 introduces the IFACD model. In Section 3, we introduce the conditional maGH distribution with higher moments dynamics. Key features such as the conditional higher comoment tensors are presented in Section 4, where we also propose a portfolio conditional density representation. The estimation of the model and the ICA algorithm are outlined in Section 5, where we report the results of a risk and portfolio management application, comparing IFACD to CHICAGO and DCC(T) models, using a representative dataset of international equity indices spanning the last two decades. Section 6 concludes. In Appendices A and B, we details the GH density and its characteristic function.

## 2 THE INDEPENDENT FACTOR MODEL

Factor ARCH models, originally introduced by Engle et al. (1990) and with foundations in the Arbitrage Pricing Theory of Ross (1976), are based on the assumption that returns are generated by a set of unobserved underlying factors that are conditionally heteroscedastic. The dependence framework is static as a consequence of large scale estimation in a multivariate setting. Consider a set of  $N$  assets whose returns  $\mathbf{r}_t$  are observed for  $T$  periods, with conditional mean  $E[\mathbf{r}_t|\mathfrak{F}_{t-1}] = \mathbf{m}_t$ , where  $\mathfrak{F}_{t-1}$  is the  $\sigma$ -field generated by the past realizations of  $\mathbf{r}_t$ , i.e.  $\mathfrak{F}_{t-1} = \sigma(\mathbf{r}_{t-1}, \mathbf{r}_{t-2}, \dots)$ . The Generalized Orthogonal GARCH (GO-GARCH) model of van der Weide (2002) maps  $\mathbf{r}_t - \mathbf{m}_t$  onto a set of unobserved independent factor  $\mathbf{f}_t$  (or "structural errors")

$$\mathbf{r}_t = \mathbf{m}_t + \boldsymbol{\epsilon}_t \quad t = 1, \dots, T \tag{1}$$

$$\boldsymbol{\epsilon}_t = \mathbf{A}\mathbf{f}_t \tag{2}$$

*The IFACD model*

where  $\mathbf{A}$  is invertible and constant over time and may be decomposed into the de-whitening  $\Sigma^{1/2}$ , representing the square root of the unconditional covariance matrix, and orthogonal matrix,  $\mathbf{U}$

$$\mathbf{A} = \Sigma^{1/2}\mathbf{U}, \quad (3)$$

and  $\mathbf{f}_t = (f_{1t}, \dots, f_{Nt})'$ . In this model it is assumed that the factors have the following specification

$$\mathbf{f}_t = \mathbf{H}_t^{1/2}\mathbf{z}_t \quad (4)$$

where  $\mathbf{H}_t = E[\mathbf{f}_t\mathbf{f}_t'|\mathfrak{F}_{t-1}]$  is a diagonal matrix with elements  $(h_{1t}, \dots, h_{Nt})$  which are the conditional variances of the factors, and  $\mathbf{z}_t = (z_{1t}, \dots, z_{Nt})'$ . The random variable  $z_{it}$  is independent of  $z_{jt-s} \forall j \neq i$  and  $\forall s$ , with  $E[z_{it}|\mathfrak{F}_{t-1}] = 0$  and  $E[z_{it}^2] = 1$ , this implies that  $E[\mathbf{f}_t|\mathfrak{F}_{t-1}] = \mathbf{0}$  and  $E[\boldsymbol{\epsilon}_t|\mathfrak{F}_{t-1}] = \mathbf{0}$ . The factor conditional variances,  $h_{i,t}$ , can be modelled as a GARCH-type process. The unconditional distribution of the factors is characterized by

$$E[\mathbf{f}_t] = \mathbf{0} \quad E[\mathbf{f}_t\mathbf{f}_t'] = \mathbf{I}_N \quad (5)$$

which, in turn, implies that

$$E[\boldsymbol{\epsilon}_t] = \mathbf{0} \quad E[\boldsymbol{\epsilon}_t\boldsymbol{\epsilon}_t'] = \mathbf{A}\mathbf{A}'. \quad (6)$$

It follows that the returns can be expressed as

$$\mathbf{r}_t = \mathbf{m}_t + \mathbf{A}\mathbf{H}_t^{1/2}\mathbf{z}_t. \quad (7)$$

The conditional covariance matrix,  $\Sigma_t \equiv E[(\mathbf{r}_t - \mathbf{m}_t)(\mathbf{r}_t - \mathbf{m}_t)'|\mathfrak{F}_{t-1}]$  of the returns is given by

$$\Sigma_t = \mathbf{A}\mathbf{H}_t\mathbf{A}'. \quad (8)$$

The estimation of GO-GARCH by maximum likelihood suffers severely from dimensionality issues. Alternative approaches such as nonlinear least squares and method of moments for the estimation of  $\mathbf{U}$  have been proposed in Boswijk and van der Weide (2006, 2011). In this paper, we estimate the  $\mathbf{U}$  by ICA as in Broda and Paoletta (2009) and Zhang and Chan (2009). One of the computational advantages offered by the generalized orthogonal approach is that following the estimation of the independent

factors, the dynamics of the marginal density parameters of those factors may be estimated separately. In this context, we propose to extend the dynamics to the full conditional density parameters to model in a multivariate setting time-varying higher moments. We consider the dynamics of the independent factors in the context of an expanded GO-GARCH model with dynamics for the full conditional parameters. While any multivariate distribution, admitting an affine representation may be used in this setup, we choose the GH distribution of Barndorff-Nielsen (1977), in the multivariate affine representation introduced by Schmidt et al. (2006), for its flexibility and rich parametrization, capturing some of the most important features of observed returns such as asymmetry and fat tails.

### 3 AUTOREGRESSIVE CONDITIONAL GH DISTRIBUTION

The GH distribution is a variance-mean mixture of normal and Generalized Inverse Gaussian (GIG) distributions. It is a flexible distribution, allowing for skewness and fat tails, nesting a large number of distributions such as the Hyperbolic, Normal Inverse Gaussian, Variance Gamma, skew-Laplace, and as limiting cases, the Normal and skew-T distributions. The  $n$ -dimensional GH distribution allows for different marginal skewness. However, Schmidt et al. (2006) point out that the margins of a random vector that is GH distributed are not mutually independent for some choice of the scaling matrix. As a consequence, they propose an alternative, non-elliptical, maGH distribution with independent margins allowed to take separate values for skewness and shape. See also Ferreira and Steel (2006) for a multivariate skew-T density with independent margins.

Based on the parametrization of Schmidt et al. (2006), the vector of returns  $\mathbf{r}_t$ , which is expressed as a linear transformation of independent factors  $\mathbf{f}_t \in \mathbb{R}^N$  as in (7), is conditionally maGH distributed

$$\mathbf{r}_t | \mathfrak{F}_{t-1} \sim maGH_N(\mathbf{m}_t, \mathbf{\Sigma}_t, \boldsymbol{\omega}_t), \quad (9)$$

where  $\boldsymbol{\omega}_t = (\omega_{1t}, \dots, \omega_{Nt})'$  and  $\omega_{it} = (\lambda_i, \alpha_{it}, \beta_{it})'$  represent the time-varying conditional shape and skew parameter vectors, respectively. In the IFACD model we assume that the standardized random variables  $z_{it}$  are conditionally distributed as a standardized GH. In general, the GH density of the random variable  $Y$  is given by

$$gh(y; \lambda, \mu, \delta, \alpha, \beta, ) = \frac{K_{\lambda-1/2} \left( \alpha \sqrt{\delta^2 + (y - \mu)^2} \right) e^{\beta(y - \mu)}}{c(\delta^2 + (y - \mu)^2)^{(1/2-\lambda)/2}}, \quad (10)$$

where

$$c = \frac{(\alpha^2 - \beta^2)^{\lambda/2}}{\sqrt{2\pi}\alpha^{\lambda-1/2}\delta^\lambda K_\lambda(\delta\sqrt{\alpha^2 - \beta^2})}.$$

Alternative distributions are nested in (10), e.g. the NIG distribution is obtained by setting  $\lambda = -\frac{1}{2}$ , the Hyperbolic with  $\lambda = \frac{\nu+1}{2}$ , and the skew-Student by setting  $\lambda = -\frac{\nu}{2}$  (with  $\nu$  representing the degrees of freedom) and  $\alpha \rightarrow |\beta|$ , where parameters  $\beta$  and  $\alpha$  are the skewn and shape parameters of the distribution, respectively. A number of location and scale invariant parameterizations of the GH distribution have been proposed in the literature. For the standardized GH distribution of  $z_{it}$  we choose the location scale invariant representation  $(\rho, \zeta)$  because we are interested in ACD modelling of skewness and kurtosis,

$$\zeta = \delta\sqrt{\alpha^2 - \beta^2}, \quad \rho = \frac{\beta}{\alpha}. \quad (11)$$

The relation between the parameters  $(\rho, \zeta)$  and  $(\alpha, \beta, \delta, \mu)$  and the standardization of the GH random variable are detailed in Appendix 6. The skew and shape parameters  $(\rho, \zeta)$  jointly determine the skewness and kurtosis (Appendix 6 reports the expressions of skewness and kurtosis of the NIG distribution). Bläesild (1981) proved that a linear transformation such as  $aX + b$  of a random variable  $X$  following a GH distribution in (10) is GH distributed with parameters  $\lambda^* = \lambda$ ,  $\alpha^* = \alpha/|a|$ ,  $\beta^* = \beta/|a|$ ,  $\delta^* = \delta|a|$ , and  $\mu^* = a\mu + b$ . The choice of alternative parameterizations involves the appropriate transformation. Given (4), the single factors,  $f_{it}, i = 1, \dots, N$ , are conditionally distributed as a  $gh(f_{it}; \lambda_i, \mu_{it}\sqrt{h_{it}}, \delta_{it}\sqrt{h_{it}}, \alpha_{it}/\sqrt{h_{it}}, \beta_{it}/\sqrt{h_{it}})$ .

We now turn to the specification of the time-varying skew and shape parameters. We adopt a quadratic dynamics for the skew parameter  $(\check{\rho}_{i,t})$  and a piece-wise linear dynamics for the shape parameter  $(\check{\zeta}_{i,t})$

$$\begin{aligned} \check{\rho}_{it} &= \chi_{0i} + \chi_{1i}\kappa_{1i}z_{it-1} + \chi_{2i}z_{it-1}^2 + \xi_{1i}\check{\rho}_{it-1} \\ \check{\zeta}_{it} &= \kappa_{0i} + \kappa_{1i}z_{it-1}\mathbf{1}_{[z_{it-1} < -1]} + \kappa_{2i}z_{it-1}\mathbf{1}_{[z_{it-1} > 1]} + \psi_{1i}\check{\zeta}_{it-1}, \end{aligned} \quad (12)$$

where  $\mathbf{1}$  is the indicator function such that positive (negative) standardized innovations, larger (smaller) than one standard deviation, have a different impact on the skew dynamics. In this way the shape responds only to shocks larger (or smaller) than one conditional standard deviation, because shocks below this threshold are less likely to be relevant and could introduce excess noise. The logistic



transform is then used to map the unconstrained processes  $\check{\rho}_{i,t}$  and  $\check{\zeta}_{i,t}$  into  $\rho_{i,t}$  and  $\zeta_{i,t}$ :

$$\rho_{it} = -0.99 + \frac{1.98}{1 + e^{-\check{\rho}_{it}}} \quad (13)$$

$$\zeta_{it} = 0.1 + \frac{24.9}{1 + e^{-\check{\zeta}_{it}}} \quad (14)$$

where the bounds of the distributional parameters are  $[-0.99, 0.99]$  and  $[0.1, 25]$  for  $\rho$  and  $\zeta$ , respectively. We limit the upper bound of  $\zeta$  to 25, since values beyond this point lead to very little change in the skewness and kurtosis, with the range 0.1 to 25 representing most of the cases, whereas the bounds for  $\rho$  are simply dictated by the GH distribution. In theory, the GIG shape parameter  $\lambda_i$  is allowed to vary for each factor, but this introduces an added layer of complexity since different combinations of  $\lambda$ ,  $\rho$  and  $\zeta$  lead to the same or close likelihood. Alternatively, choosing a value of  $\lambda$  equal to  $-0.5$ , which corresponds to the NIG distribution, we have enough flexibility to account for the observed non-Gaussian features of financial time series. This is our choice in the empirical implementation of IFACD model in Section 5.

In the next section, we present the conditional co-moments and portfolio conditional density implied by the IFACD model, employed in the approximation of the expected utility in Section 5.4.

## 4 CONDITIONAL CO-MOMENTS AND PORTFOLIO CONDITIONAL DENSITY

The novelty of the IFACD model is that the co-moments are now time-varying, as a consequence of the ACD specification of the conditional density of the standardized innovations. The conditional co-moments of  $\mathbf{r}_t$  of order 3 and 4 are represented as tensor matrices

$$\mathbf{M}_t^3 = \mathbf{A} \mathbf{M}_{f,t}^3 (\mathbf{A} \otimes \mathbf{A})', \quad \mathbf{M}_t^4 = \mathbf{A} \mathbf{M}_{f,t}^4 (\mathbf{A} \otimes \mathbf{A} \otimes \mathbf{A})', \quad (15)$$

where  $\mathbf{M}_{f,t}^3$  and  $\mathbf{M}_{f,t}^4$  are the  $(N \times N^2)$  conditional third comoment matrix and the  $(N \times N^3)$  conditional fourth comoment matrix of the factors, respectively.  $\mathbf{M}_{f,t}^3$  and  $\mathbf{M}_{f,t}^4$ , are defined as

$$\mathbf{M}_{f,t}^3 = \begin{bmatrix} \mathbf{M}_{1,f,t}^3, \mathbf{M}_{2,f,t}^3, \dots, \mathbf{M}_{N,f,t}^3 \end{bmatrix} \quad (16)$$

$$\mathbf{M}_{f,t}^4 = \begin{bmatrix} \mathbf{M}_{11,f,t}^4, \mathbf{M}_{12,f,t}^4, \dots, \mathbf{M}_{1N,f,t}^4 | \dots | \mathbf{M}_{N1,f,t}^4, \mathbf{M}_{N2,f,t}^4, \dots, \mathbf{M}_{NN,f,t}^4 \end{bmatrix} \quad (17)$$

*The IFACD model*

where  $\mathbf{M}_{k,f,t}^3, k = 1, \dots, N$  and  $\mathbf{M}_{kl,f,t}^4, k, l = 1, \dots, N$  are the  $(N \times N)$  submatrices of  $\mathbf{M}_{f,t}^3$  and  $\mathbf{M}_{f,t}^4$ , respectively, with elements

$$\begin{aligned} m_{ijk,f,t}^3 &= E[f_{i,t}f_{j,t}f_{k,t}|\mathfrak{F}_{t-1}] \\ m_{ijkl,f,t}^4 &= E[f_{i,t}f_{j,t}f_{k,t}f_{l,t}|\mathfrak{F}_{t-1}]. \end{aligned}$$

Since the factors  $f_{it}$  can be decomposed as  $z_{it}\sqrt{h_{it}}$ , and given the assumptions on  $z_{it}$ , then  $E[f_{i,t}f_{j,t}f_{k,t}|\mathfrak{F}_{t-1}] = 0$ . It is also true that for  $i \neq j \neq k \neq l$ ,  $E[f_{i,t}f_{j,t}f_{k,t}f_{l,t}|\mathfrak{F}_{t-1}] = 0$  and when  $i = j$  and  $k = l$

$$E[f_{i,t}f_{j,t}f_{k,t}f_{l,t}|\mathfrak{F}_{t-1}] = h_{it}h_{kt}.$$

Thus, under the assumption of mutual independence, all elements in the conditional co-moments matrices with at least 3 different indices are zero. Finally, we standardize the conditional co-moments to obtain conditional coskewness and cokurtosis of  $\mathbf{r}_t$

$$\mathbf{S}_{ijk,t} = \frac{m_{ijk,t}^3}{(\sigma_{i,t}\sigma_{j,t}\sigma_{k,t})}, \quad \mathbf{K}_{ijkl,t} = \frac{m_{ijkl,t}^4}{(\sigma_{i,t}\sigma_{j,t}\sigma_{k,t}\sigma_{l,t})}, \quad (18)$$

where  $\mathbf{S}_{ijk,t}$  represents the coskewness between elements  $i, j, k$  of  $\mathbf{r}_t$ ,  $\sigma_{i,t}$  the standard deviation of  $\mathbf{r}_{i,t}$ , and in the case of  $i = j = k$  represents the skewness of asset  $i$  at time  $t$ , and similarly for the cokurtosis tensor  $\mathbf{K}_{ijkl,t}$ .<sup>1</sup>

An important question that can be addressed in this framework is the determination of the portfolio conditional density, an issue of vital importance in the risk management application. Let  $R_t$  be the portfolio return

$$R_t = \mathbf{w}'_t \mathbf{r}_t = \mathbf{w}'_t \mathbf{m}_t + (\mathbf{w}'_t \mathbf{A} \mathbf{H}_t^{1/2}) \mathbf{z}_t \quad (19)$$

where  $\mathbf{H}_t^{1/2}$  is obtained from the ACD dynamics of estimated  $\mathbf{f}_t$ . The portfolio conditional variance,

---

<sup>1</sup>A natural application of return co-moments matrices is the news impact surface (as first suggested by Kroner and Ng, 1998, and extended by Jondeau and Rockinger, 2009) which provides additional insights on the effect of factor shocks on the asset co-moments. The results of this application are available upon request.

skewness and kurtosis in closed form are

$$\begin{aligned}\sigma_{R,t}^2 &= \mathbf{w}'_t \boldsymbol{\Sigma}_t \mathbf{w}_t, \\ s_{R,t} &= \frac{\mathbf{w}'_t \mathbf{M}_t^3 (\mathbf{w}_t \otimes \mathbf{w}_t)}{(\mathbf{w}'_t \boldsymbol{\Sigma}_t \mathbf{w}_t)^{3/2}}, \\ k_{R,t} &= \frac{\mathbf{w}'_t \mathbf{M}_t^4 (\mathbf{w}_t \otimes \mathbf{w}_t \otimes \mathbf{w}_t)}{(\mathbf{w}'_t \boldsymbol{\Sigma}_t \mathbf{w}_t)^2},\end{aligned}\tag{20}$$

where  $\boldsymbol{\Sigma}_t$ ,  $\mathbf{M}_t^3$  and  $\mathbf{M}_t^4$  are derived in (8) and (15), respectively. The portfolio conditional density may be obtained via the inversion of the characteristic function through the FFT method as in Chen et al. (2007) (see Appendix 6 for details) or by simulation. We implement the FFT for its accuracy and speed. While the  $N$ -dimensional NIG distribution is closed under convolution, when the distributional parameters  $\alpha$  and  $\beta$  are allowed to be different across assets, as in the case of IFACD model, this property no longer holds. Provided that  $\mathbf{z}_t$  is a  $N$ -dimensional vector of innovations, marginally distributed as 1-dimensional standardized GH, the density of weighted asset return,  $w_{it}r_{it}$ , is

$$w_{it}r_{it} = (w_{i,t}m_{i,t} + \bar{w}_{i,t}z_{i,t}) \sim maGH_1 \left( \bar{w}_{i,t}\mu_{i,t} + w_{i,t}m_{i,t}, |\bar{w}_{i,t}| \delta_{i,t}, \lambda_i, \frac{\alpha_{i,t}}{|\bar{w}_{i,t}|}, \frac{\beta_{i,t}}{|\bar{w}_{i,t}|} \right)\tag{21}$$

where  $\bar{\mathbf{w}}'_t$  is equal to  $\mathbf{w}'_t \mathbf{A} \mathbf{H}_t^{1/2}$ , and  $\bar{w}_{i,t}$  is the  $i$ -th element of  $\bar{\mathbf{w}}_t$ ,  $m_{i,t}$  the conditional mean of the  $i$ -th underlying asset. In order to obtain the density of the portfolio, we must sum the individual weighted densities of  $z_{i,t}$ . The characteristic function of the portfolio return  $R_t$  is

$$\varphi_R(u) = \prod_{i=1}^n \varphi_{\bar{w}Z_i}(u) = \exp \left( iu \sum_{j=1}^d \bar{\mu}_j + \sum_{j=1}^d \left( \frac{\lambda_j}{2} \log \left( \frac{\gamma}{v} \right) + \log \left( \frac{K_{\lambda_j}(\bar{\delta}_j \sqrt{v})}{K_{\lambda_j}(\bar{\delta}_j \sqrt{\gamma})} \right) \right) \right)\tag{22}$$

where,  $\gamma = \bar{\alpha}_j^2 - \bar{\beta}_j^2$ ,  $v = \bar{\alpha}_j^2 - (\bar{\beta}_j + iu)^2$ , and  $(\bar{\alpha}_j, \bar{\beta}_j, \bar{\delta}_j, \bar{\mu}_j)$  are the scaled versions of the parameters  $(\alpha, \beta_i, \delta_i, \mu_i)$  as shown in (21). The density is accurately approximated by FFT as follows

$$f_R(r) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{(-iur)} \varphi_R(u) du \approx \frac{1}{2\pi} \int_{-s}^s e^{(-iur)} \varphi_R(u) du.\tag{23}$$

Expression (23) is the base for the calculation of VaR in the empirical application reported in Section 5.

## 5 EMPIRICAL APPLICATION

In this section, we report results of an empirical exercise using data on 14 MSCI Global Equity indices representing a cross section of countries in North America (U.S., Canada, Mexico), Asia (Australia, Hong Kong, Japan, Singapore) and Europe (Germany, France, Spain, Italy, U.K., Switzerland, Sweden), from 12/08/1996 to 28/12/2011, and obtained from Yahoo Finance. The period includes the Asian Financial Crisis of 1997, the Russian Financial Crisis of 1998, the Dot-Com bubble of 2000 and subsequent economic downturn of 2002, the Chinese Stock Bubble of 2007, the US Bear Market of 2007-2009, the European sovereign debt crisis of 2010 as well as the flash crash of May 2010.<sup>2</sup> First, we report (Section 5.2) an in-sample comparison of the IFACD and CHICAGO models in order to obtain some insight into the types of dynamics and parametrization of the factors. In Section 5.3, we report the results of an out-of-sample risk forecast evaluation using randomly weighted portfolios to avoid any bias arising from the uncertainty in selecting any particular set of weights. In Section 5.4, a Taylor series expansion of the Constant Absolute Risk Aversion (CARA) utility function is used for an out-of-sample portfolio allocation to contrast the time-varying versus the static representation of higher conditional co-moment matrices.

In the following subsection, we briefly describe the ICA method, used to estimate the independent factors, and we introduce the likelihood function.

### 5.1 Estimation Procedure

The estimation procedure of the IFACD model can be summarized as follows. First, we compute the ICA of the data  $\mathbf{z}_t = \widehat{\Sigma}^{-1/2} \widehat{\boldsymbol{\epsilon}}_t$ , where  $\widehat{\boldsymbol{\epsilon}}_t$  are the OLS residuals, i.e.  $\widehat{\boldsymbol{\epsilon}}_t = \mathbf{r}_t - \widehat{\mathbf{m}}_t$ , and  $\widehat{\Sigma}^{1/2}$  is obtained from the eigenvalue decomposition of the OLS residual covariance matrix. ICA provides an estimate of the orthogonal matrix  $\mathbf{U}$  in (3) as in Broda and Paoletta (2009) and Zhang and Chan (2009). The ICA is a computational method for separating multivariate mixed signals,  $\mathbf{x} = [x_1, \dots, x_n]'$ , into additive statistically independent and non-Gaussian components,  $\mathbf{s} = [s_1, \dots, s_n]'$ , such that  $\mathbf{x} = \mathbf{B}\mathbf{s}$ . The estimate of the linear mixing matrix  $\mathbf{B}$  can be obtained via the algorithm, proposed and implemented by Hyvärinen and Oja (2000). The FastICA algorithm is based on negentropy, which is an optimal estimate of non-Gaussianity, as shown by Comon (1994), is invariant to invertible linear transformations. Thus the estimated  $\mathbf{f}_t$  are obtained as  $\hat{\mathbf{f}}_t = \widehat{\Sigma}^{-1/2} \widehat{\mathbf{U}} \mathbf{z}_t$ . Second, because of the assumption of indepen-

---

<sup>2</sup>The tickers of these iShare tradeable indices, in the order presented are: SPY, EWC, EWW, EWA, EWH, EWJ, EWS, EWG, EWQ, EWP, EWI, EWU, EWL, EWD.

dence, the likelihood function of the IFACD model is greatly simplified. The conditional log-likelihood function is expressed as the sum of the individual conditional log-likelihoods, derived from the conditional marginal densities of the factors, i.e.,  $gh(\hat{f}_{it}) \equiv gh(\hat{f}_{it}; \lambda_i, \mu_{it}\sqrt{h_{it}}, \delta_{it}\sqrt{h_{it}}, \alpha_{it}/\sqrt{h_{it}}, \beta_{it}/\sqrt{h_{it}})$ , plus a term for the mixing matrix  $\mathbf{A}$ , estimated in the first step by FastICA

$$L(\hat{\epsilon}_t | \boldsymbol{\theta}, \mathbf{A}) = T \log |\mathbf{A}^{-1}| + \sum_{t=1}^T \sum_{i=1}^N \log \left( gh(\hat{f}_{it} | \theta_i) \right) \quad (24)$$

where  $\boldsymbol{\theta}$  is the vector of unknown parameters in the marginal densities. The possibility of modelling the independent factors separately not only increases the flexibility of the model but also its computational feasibility, since the multivariate estimation reduces to  $N$  univariate optimization steps plus a term which depends on the factor loading matrix.

## 5.2 In-sample Estimation

In Table 1, we report the results of the first in-sample fit of the IFACD and CHICAGO models for the period 19/03/1996 to 17/03/2000, in order to illustrate the factors' dynamics and their significance. The CHICAGO model consists of (1)-(8), but with the conditional distribution of  $f_{it}$  which has only the second moment time-varying and constant higher moments, and the matrix  $\mathbf{U}$  in (3) estimated by FastICA. Both models are estimated under the maNIG distribution. Because of the presence of first degree autocorrelation in the MSCI index dataset, the returns were demeaned and filtered with an AR(1) model prior to applying the FastICA algorithm. In both models, the factor variance dynamics follow a GARCH(1,1) model with parameters  $(\omega, \alpha_1, \beta_1)$ , and in the IFACD the skew and shape dynamics follow a first order quadratic and piece-wise linear model with parameters  $(\chi_0, \chi_1, \chi_2, \xi_1)$  and  $(\kappa_0, \kappa_1, \kappa_2, \psi_1)$  respectively, as in (12). Given that ICA is conceived as a linear noiseless model, the standard errors are computed only in the estimation of the factors' conditional distribution parameters.

Starting with the CHICAGO model, the very small absolute value and lack of significance of the skew parameter  $\rho$  indicates that skewness is not very pronounced in this sample, and the skew dynamics of the IFACD model somewhat bear this out since very few factors appear to have significant skew dynamics. The same cannot be said of the shape parameter  $\zeta$  which is significant in the majority of factors and varies between 1.7 and 2.5, with the exception of Factor 10 which does not appear to have significant skew or shape, and in combination with a zero skew parameter translates to an excess kurtosis of between 1.7 and 1.2 respectively. The shape dynamics are also significant for the

majority of the factors in the IFACD model. With regard to the variance dynamics, they appear to be the same across the two models, though there is some evidence from Factors 1, 12 and 14 that when persistence is very high in the CHICAGO model, the IFACD model persistence instead is much lower as the shape dynamics accommodate expansions and contractions in the conditional density shape which would otherwise be completely captured by the GARCH variance dynamics. Finally, while the factor log-likelihoods (and hence model log-likelihood) of the IFACD model are always higher than those of the CHICAGO model, the higher BIC of the former indicates that there is a small penalty for the overparametrization.

### 5.3 Out-of-sample Risk Forecast

The out-of-sample comparison is based on one-step ahead forecasts. We estimate the models every five days, keeping a constant sample size, for a total of 522 re-estimations resulting in 2610 forecasts. Starting from 17/03/2000, the last 4 years of daily log returns are used to estimate IFACD models, CHICAGO, and DCC(T). In order to evaluate the contribution that the ACD specification gives to the model's performance, in addition to the IFACD model with the time-varying  $\rho_t$  and  $\zeta_t$ , we consider two restricted versions of the model which have constant skew and time-varying shape, i.e.  $(\rho, \zeta_t)$ , and time-varying skew and constant shape, i.e.  $(\rho_t, \zeta)$ , respectively. We also include the DCC(T) model so as to gauge the cost of IFACD which is based on unconditional independence. The model is specified and estimated as in Bauwens and Laurent (2005).

The consistency between the IFACD and CHICAGO models was guaranteed by using the same mixing matrix  $\mathbf{A}$  across the two models for each estimation window so that the only difference is purely in the dynamics of the factors. Because of the non-dynamic nature of the mixing matrix in the IFACD model, the rolling window re-estimation scheme with a fixed size of 4 years enables to capture any change to the loadings, though tracking such changes is not trivial since the independent components are identified only up to a permutation and scaling of the sources. To evaluate the forecast performance of the models, a weighted linear combination of the forecast density was used in order to form portfolios from which measures could easily be calculated. To avoid bias from any particular weighting scheme, 1000 + 1 random portfolios are generated by sampling weights from the exponential distribution (the 1001<sup>th</sup> is the equally weighted portfolio) and dividing by the sum of the randomly generated deviates to create the full investment constraint. The weighted densities of the IFACD and CHICAGO models are estimated using the FFT method described in Section 4 from which quantile and distribution

functions are easily created. For DCC(T) model the portfolio moments used to compute functions of the density are obtained using the standard linear and quadratic transformations of the forecast mean and covariance matrix. The following tests are used to evaluate the forecasting accuracy of the risk models: Berkowitz (2001) for testing the predictive density, Kupiec (1995) and Christoffersen (1998) for VaR exceedances, and Christoffersen and Pelletier (2004) for VaR Durations.

Table 2 reports the result under the different tests for the equally weighted (EW) and average of the randomly weighted (RAND) portfolios. For the Berkowitz test, there is no significant difference between the factor models, with a rejection rate among the randomly weighted portfolios of about 16% and 14% for the IFACD( $\rho_t, \zeta_t$ ) and CHICAGO models respectively, indicating that overall both models fit the out-of-sample forecast density well. The same is true for the IFACD models with constrained ACD dynamics. The DCC(T) model on the other hand does not fit the conditional weighted forecast densities very well, with more than 50% rejection rate.

For the VaR tests, both at the 1% and 5% coverage rates, the IFACD( $\rho_t, \zeta_t$ ) model performs better than the other two IFACD models and the CHICAGO, as evidenced by the large difference in rejection rates. In the VaR Duration test with 1% coverage, the IFACD( $\rho_t, \zeta_t$ ) and IFACD( $\rho, \zeta_t$ ) dominate the CHICAGO. However, the DCC(T) outperforms the factor models and this is indicative of the value of dynamic correlation since the duration indirectly tests for clustering of tail events which is not likely to be fully filtered out in a static independent factor framework.

#### 5.4 Out-of-sample Optimal Portfolio Forecast

We consider an alternative approach to the standard quadratic model of Markowitz (1952) is based on the approximation of the utility function via a Taylor series expansion. In each period, an investor allocates capital to maximize the conditional expected utility,  $E_{t-1}[U(W_t)]$ , over the next period wealth  $W_t$ , with initial wealth equal to 1. The optimal allocation problem may be formulated as:

$$\max_{\mathbf{w}_t} E_{t-1} [U(W_t)], \quad \text{s.t.} \quad \sum_{i=1}^N w_{ti} = 1, \quad w_{ti} \geq 0 \quad \forall i, \quad (25)$$

where we assume the absence of a riskless asset, so that the sum of the weights  $w_{ti}$  sums to one; we also exclude the possibility of short-selling. The expected utility can be written, using a Taylor series expansion (see for analogous approach Jurczenko and Maillet (2006) and Jondeau and Rockinger

(2006)), as

$$E_{t-1} [U(W)] = U(\bar{W}_t) + U^{(1)}(\bar{W}_t) E_{t-1} [W_t - \bar{W}_t] + \frac{1}{2} U^{(2)}(\bar{W}_t) E_{t-1} [(W_t - \bar{W}_t)^2] + \frac{1}{3} U^{(3)}(\bar{W}_t) E_{t-1} [(W_t - \bar{W}_t)^3] + \frac{1}{4} U^{(4)}(\bar{W}_t) E_{t-1} [(W_t - \bar{W}_t)^4] + O(W_t^4), \quad (26)$$

where  $\bar{W}_t = \mathbf{w}'_t E_{t-1} [\mathbf{r}_t]$  is the expected portfolio return and  $O(W_t^4)$  represents the remainder of the Taylor series due to the truncation.

Note that this approximation includes two moments more than the mean-variance criterion, with the skewness and kurtosis are directly related to investor preferences (dislike) for odd (even) moments, under certain mild assumptions given in Scott and Horvath (1980). The expected utility can then be approximated as

$$E_{t-1} [U(W_t)] \approx U(\bar{W}_t) + \frac{1}{2} U^{(2)}(\bar{W}_t) \sigma_{R,t}^2 + \frac{1}{3!} U^{(3)}(\bar{W}_t) \tilde{s}_{R,t} + \frac{1}{4!} U^{(4)}(\bar{W}_t) \tilde{k}_{R,t}. \quad (27)$$

where  $\tilde{s}_{R,t} \equiv E_{t-1} [(W_t - \bar{W}_t)^3]$  and  $\tilde{k}_{R,t} \equiv E_{t-1} [(W_t - \bar{W}_t)^4]$  are the nonstandardized versions of conditional skewness and kurtosis, namely the numerators in (20). When the CARA utility function is used, then the approximation resolves to:

$$E_{t-1} [U(W_t)] = E_{t-1} [-\exp(-\eta W_t)] \approx -\exp(-\eta \bar{W}_t) \left[ 1 + \frac{\eta}{2} \sigma_{R,t}^2 - \frac{\eta^3}{3!} \tilde{s}_{R,t} + \frac{\eta^4}{4!} \tilde{k}_{R,t} \right], \quad (28)$$

where  $\eta$  represents the investor's constant absolute risk aversion, with higher (lower) values representing higher (lower) aversion. It is evident from (28) that the conditional kurtosis has a negative impact on the expected utility while the positive conditional skewness has a positive one. We maximize this function for all rolling forecasts using a nonlinear Sequential Quadratic Programming solver and making use of the first order derivatives given in Jondeau and Rockinger (2006). Upper bounds (50%) are imposed on the assets weights.

Table 3 shows the results for the IFACD, CHICAGO and DCC models of CARA utility portfolios maximized under 4 different risk aversion coefficients from the mildly risk averse to extremely risk averse investor. The results of this exercise confirm the benefit of considering higher conditional moments in portfolio allocation. As the risk aversion coefficient increases, and consequently more weight is given to the higher moments in the conditional expected utility, the IFACD( $\rho_t, \zeta_t$ ) model progressively outperforms the CHICAGO model. This is immediately evident from the  $p$ -value of the



model confidence set (MCS) of Hansen et al. (2011), using as loss function the negative of the portfolio returns, which overwhelmingly rejects both the CHICAGO and DCC(T) models from  $\eta = 10$  onwards, with 90% confidence.

Turning to IFACD models with constrained ACD dynamics, we have very similar results to those of IFACD( $\rho_t, \zeta_t$ ). This can be explained by the fact that both the skew and shape parameters jointly determine the tail behavior in the NIG/GH distribution, so that when there is very little variation in the conditional asymmetry, as in our sample, the skew parameter is mostly adjusting to compensate for the fixed shape parameter in determining the dynamics in the tail behavior. The portfolios based on IFACD models underweight securities with high predicted kurtosis and negative skewness, and as a result, when  $\eta$  increases and more emphasis is placed on higher moments, the average return and Risk-Reward (RR) ratio, in Table 3, are significantly higher than the CHICAGO and DCC(T) models. A slow progression is seen when it comes to the significance of RR differences given by the test of Ledoit and Wolf (2008), with the IFACD( $\rho_t, \zeta_t$ ) model RR ratio significantly better at  $\eta = 25$  with 90% confidence. Not surprisingly, the DCC model based on the CARA utility expansion with only 3 moments fares worse at all levels of risk aversion.

## 6 CONCLUSIONS

In portfolio and risk management analysis, it is important to consider not only asymmetric and heavy tailed multivariate distributions but also time-varying skewness and kurtosis. Modelling the conditional density dynamics in a multivariate setup has proved unfeasible mainly because of the difficulty in dealing with tractable representations of non-elliptical multivariate distributions. In this paper, we propose the IFACD model where the returns have a maGH distribution with time-varying higher moments. These features allow us to model extreme behaviors of financial markets turmoils.

The IFACD model uses time-varying distributional parameters in a multivariate setting, for truly large scale models. Further, the IFACD model has closed form higher moments and semi-analytic expression for the portfolio density which have important implications for portfolio allocation and risk management analysis. Modelling higher moments as a time-varying processes is highly appropriate for periods of market stress, when GARCH models cannot accommodate extreme events. We report evidence of this via large out-of-sample risk management and portfolio applications: using 14 MSCI Global Equity indices at daily frequency, over the period 1996-2011, the IFACD model outperforms

CHICAGO and DCC(T) models.

The findings in this paper suggest further developments. For instance, it would be interesting to study consistent procedures able to provide more parsimonious independent factor representations. Further, the use of IFACD model with high-frequency data or realized measures would be a useful extension to mimic the evolution of volatilities in real time. We leave these issues to future research.

## **ACKNOWLEDGEMENT**

This work started while the second author was visiting the Centre for Econometric Analysis at Cass Business School in summer-fall 2009. We thank participants in the 4th CSDA International Conference on Computational and Financial Econometrics (London, 10-12 December 2010), the Fourth Italian Congress of Econometrics and Empirical Economics (Pisa, 19-21 January 2011). We wish to thank the Editor, Esfandiar Maasoumi, an Associate Editor and two anonymous referees for useful comments and suggestions which greatly helped to improve the paper. The usual disclaimer applies.

Table 1: IFACD and CHICAGO Models: Parameter Estimates

IFACD														
	$f_1$	$f_2$	$f_3$	$f_4$	$f_5$	$f_6$	$f_7$	$f_8$	$f_9$	$f_{10}$	$f_{11}$	$f_{12}$	$f_{13}$	$f_{14}$
$\omega$	0.228***	0.012	0.056	0.012	0.067*	0.012	0.059	0.040	0.085	0.054***	0.018***	0.203***	0.430***	0.139**
$\alpha_1$	0.090***	0.055***	0.087	0.021	0.079***	0.017	0.114	0.067	0.064	0.054***	0.037***	0.071	0.290	0.038***
$\beta_1$	0.691***	0.935***	0.869***	0.969	0.850***	0.972**	0.832***	0.888***	0.846***	0.898***	0.946***	0.731***	0.350**	0.820***
$\chi_0$	0.077**	-0.090	0.029	0.284	-0.277**	0.218	0.031	-0.230	0.028**	-0.025	-0.036	0.061	0.056	-0.173
$\chi_1$	0.145**	0.133	0.234***	-0.286	0.008	0.050	0.125	0.303*	0.042	0.783**	0.193	0.109	0.227*	0.568
$\chi_2$	-0.068*	-0.039	-0.061	-0.206	0.071*	-0.178	-0.018	-0.071*	-0.025**	-0.059	0.092	-0.020	0.012	0.382
$\xi_1$	0.921***	0.210	0.719	0.002	0.000	0.009	0.895***	0.392*	0.929***	0.438**	0.007	0.000	0.585	0.246
$\kappa_0$	-2.373	-2.408***	-0.135	-1.096	-0.255*	-2.317	-0.015	-0.010	-2.791***	0.320	-1.614*	0.246	-2.273	-2.365***
$\kappa_1$	0.489**	0.244	0.223***	0.449	-0.392**	0.453	0.380	-0.052	0.479	0.174	-0.746	0.572*	0.681	-0.247
$\kappa_2$	-0.943***	-0.045	-0.633***	-0.474	0.089	-0.289	-0.512***	-0.132	0.955*	-1.000	0.981	-0.999	-0.998***	1.000
$\psi_1$	0.001	0.000	0.882***	0.425	0.941***	0.000	0.911***	0.986***	0.008	0.957***	0.403	0.889***	0.001	0.000
$\alpha_1 + \beta_1$	0.781	0.990	0.956	0.989	0.929	0.989	0.946	0.955	0.911	0.951	0.983	0.802	0.641	0.858
<i>Factor</i> <sub>LL</sub>	-1391.15	-1356.72	-1362.39	-1400.23	-1353.69	-1397.26	-1327.67	-1342.91	-1346.19	-1418.20	-1384.97	-1416.16	-1373.69	-1383.06
<i>Factor</i> <sub>BIC</sub>	2.83	2.76	2.78	2.85	2.76	2.85	2.71	2.74	2.74	2.89	2.82	2.88	2.80	2.82
<i>Model</i> <sub>LL</sub>	41148													
<i>Model</i> <sub>AIC</sub>	-5.80													
<i>Model</i> <sub>BIC</sub>	-5.72													
CHICAGO														
	$f_1$	$f_2$	$f_3$	$f_4$	$f_5$	$f_6$	$f_7$	$f_8$	$f_9$	$f_{10}$	$f_{11}$	$f_{12}$	$f_{13}$	$f_{14}$
$\omega$	0.044*	0.010	0.039	0.015	0.069*	0.012	0.081**	0.046**	0.063**	0.057	0.025	0.001	0.299*	0.004
$\alpha_1$	0.031*	0.050***	0.045**	0.019***	0.096***	0.018*	0.086**	0.070***	0.060***	0.052*	0.042**	0.000	0.135**	0.001
$\beta_1$	0.926***	0.942***	0.915***	0.967***	0.835***	0.971***	0.826***	0.883***	0.873***	0.893***	0.934***	0.999***	0.561**	0.995***
$\rho$	0.071	-0.088	-0.102	0.065	-0.108*	0.010	0.048	-0.098	0.031	-0.011	0.009	0.007	0.086	0.005
$\zeta$	1.781***	1.944***	1.840***	2.343***	1.781***	1.832***	1.860***	1.654	1.610***	15.000	2.573***	2.550***	1.917***	1.743**
$\alpha_1 + \beta_1$	0.957	0.992	0.960	0.986	0.931	0.986	0.912	0.952	0.933	0.944	0.976	0.999	0.696	0.996
<i>Factor</i> <sub>LL</sub>	-1400.19	-1357.95	-1372.24	-1405.83	-1357.17	-1402.68	-1333.49	-1347.28	-1349.96	-1420.38	-1387.01	-1420.38	-1378.65	-1392.05
<i>Factor</i> <sub>BIC</sub>	2.81	2.73	2.75	2.82	2.72	2.81	2.68	2.70	2.71	2.86	2.78	2.85	2.77	2.79
<i>Model</i> <sub>LL</sub>	41074													
<i>Model</i> <sub>AIC</sub>	-5.80													
<i>Model</i> <sub>BIC</sub>	-5.76													

**Note:** The Table reports the parameter estimates under the IFACD and CHICAGO models, with maNIG distribution, for the log returns of 14 MSCI indices from 19/03/1996 to 17/03/2000 (1010 days). The conditional variance of the factors follows a GARCH(1,1) model:  $h_{i,t} = \omega_j + \alpha_j f_{i,t-1} + \beta_j h_{i,t-1}$ ,  $i = 1, \dots, 14$ . The conditional skew dynamics of the factors is bounded through a logistic transformation such that  $\rho_{it} = -0.99 + \frac{1.98}{1 + \exp(-\rho_{it})}$ , where the unconstrained parameters follow a first order quadratic model:

$\check{\rho}_{it} = \chi_0 i + \chi_1 i z_{it-1} + \theta_{2i} z_{it-1}^2 + \xi_{1i} \check{\rho}_{it-1} + \zeta_{it}$ . The conditional shape dynamics of the factors is bounded through a logistic transformation such that  $\zeta_{it} = 0.1 + \frac{24.9}{1 + \exp(-\zeta_{it})}$ , where the unconstrained parameters follow first order piecewise linear model:  $\check{\zeta}_{it} = \kappa_{0i} + \kappa_{1i} z_{it-1} \mathbf{1}_{[z_{it-1} < -1]} + \kappa_{2i} z_{it-1} \mathbf{1}_{[z_{it-1} > 1]} + \psi_{1i} \check{\zeta}_{it-1}$ .

The \*, \*\* and \*\*\* next to the parameters denote significance at the 10%, 5% and 1% levels respectively. Robust standard errors of White (1982) were calculated. The individual factor volatility persistence ( $\alpha_1 + \beta_1$ ), log-likelihood and BIC are reported as is the overall model log-likelihood, AIC and BIC for comparison between the two models.

Table 2: VaR Exceedances and Density Forecast Tests

	<i>Berkowitz</i>	<i>VaR<sub>CC1%</sub></i>	<i>VaR<sub>CC5%</sub></i>	<i>VaR<sub>Dur1%</sub></i>	<i>VaR<sub>Dur5%</sub></i>
<b>IFACD</b> ( $\rho_t, \zeta_t$ )					
EW					
<i>p</i> -value	0.097	0.359	0.055	0.085	0.005
RAND					
<i>p</i> -value	0.160	0.170	0.100	0.070	0.021
% Rejection	15.6	22.8	37.9	61.9	88.7
<b>IFACD</b> ( $\rho, \zeta_t$ )					
EW					
<i>p</i> -value	0.084	0.150	0.094	0.027	0.009
RAND					
<i>p</i> -value	0.160	0.200	0.100	0.130	0.030
% Rejection	20.3	27.9	40.0	40.7	85.0
<b>IFACD</b> ( $\rho_t, \zeta$ )					
EW					
<i>p</i> -value	0.108	0.198	0.051	0.021	0.006
RAND					
<i>p</i> -value	0.170	0.130	0.090	0.030	0.010
% Rejection	13.2	38.6	45.1	84.0	93.9
<b>CHICAGO</b>					
EW					
<i>p</i> -value	0.140	0.04	0.05	0.009	0.009
RAND					
<i>p</i> -value	0.220	0.070	0.100	0.040	0.020
% Rejection	13.7	60.8	53.3	74.7	87.3
<b>DCC(T)</b>					
EW					
<i>p</i> -value	0.022	0.002	0.004	0.068	0.168
RAND					
<i>p</i> -value	0.080	0.010	0.020	0.170	0.160
% Rejection	53.8	95.5	93.0	19.9	29.9

**Note:** The Table reports the out-of-sample performance of the IFACD, CHICAGO (maNIG), and DCC(T) models for 14 MSCI indices for the period 11/08/2000 to 28/12/2010 (2610 days) based on the density test of Berkowitz (2001), the conditional coverage test for VaR exceedances of Christoffersen (1998) (*VaR<sub>CC</sub>*), and the Duration of VaR Exceedances test of Christoffersen and Pelletier (2004) (*VaR<sub>Dur</sub>*). The null hypothesis for all tests is that the models are correctly specified for which the table reports the *p*-values of the test under an equally weighted (EW) portfolio, and the average *p*-value of 1,000 random weighted (RAND) portfolios with full investment budget constraint. For the RAND portfolio, the number of rejections of the null hypothesis at the 5% level of significance is also reported.

Table 3: Time-varying Higher Co-moments Portfolio with CARA

	IFACD ( $\rho_t, \zeta_t$ )	IFACD ( $\rho, \zeta_t$ )	IFACD ( $\rho_t, \zeta$ )	CHICAGO	DCC(T)
$\eta = 1$					
$\bar{W}_T$	75.47	81.21	83.40	79.72	57.21
$\hat{\mu}$	0.0018	0.0019	0.0019	0.0018	0.0017
$\hat{\sigma}$	0.0169	0.0190	0.0190	0.0170	0.0168
RR	1.69	1.56	1.57	1.71	1.60
LW [ <i>p</i> -value]		[0.276]	[0.299]	[0.224]	[0.248]
MCS [ <i>p</i> -value]	[0.490]	[0.490]	[1.000]	[0.490]	[0.455]
log(Relative Wealth)		0.07	0.10	0.05	-0.28
$\eta = 5$					
$\bar{W}_T$	94.07	93.56	94.70	58.60	43.46
$\hat{\mu}$	0.0019	0.0019	0.0019	0.0017	0.0016
$\hat{\sigma}$	0.0175	0.0175	0.0175	0.0161	0.0159
RR	1.72	1.71	1.72	1.66	1.57
LW [ <i>p</i> -value]		[0.470]	[0.863]	[0.564]	[0.228]
MCS [ <i>p</i> -value]	[0.871]	[0.871]	[1.000]	[0.160]	[0.057]
log(Relative Wealth)		-0.01	0.01	-0.47	-0.77
$\eta = 10$					
$\bar{W}_T$	64.20	62.85	62.00	39.84	25.33
$\hat{\mu}$	0.0017	0.0017	0.0017	0.0015	0.0014
$\hat{\sigma}$	0.0163	0.0163	0.0163	0.0154	0.0151
RR	1.69	1.68	1.67	1.58	1.42
LW [ <i>p</i> -value]		[0.294]	[0.185]	[0.198]	[0.020]
MCS [ <i>p</i> -value]	[1.000]	[0.607]	[0.607]	[0.070]	[0.005]
log(Relative Wealth)		-0.02	-0.03	-0.48	-0.93
$\eta = 25$					
$\bar{W}_T$	23.83	24.73	24.36	17.82	11.88
$\hat{\mu}$	0.0013	0.0013	0.0013	0.0012	0.0010
$\hat{\sigma}$	0.0145	0.0145	0.0145	0.0142	0.0138
RR	1.44	1.46	1.45	1.34	1.20
LW [ <i>p</i> -value]		[0.121]	[0.373]	[0.107]	[0.006]
MCS [ <i>p</i> -value]	[0.246]	[1.000]	[0.693]	[0.069]	[0.004]
log(Relative Wealth)		0.04	0.02	-0.29	-0.70

**Note:** The Table reports the out-of-sample performance of the IFACD, CHICAGO (maNIG), and DCC (T) models from the optimization of the CARA utility approximation using only the first 4 co-moment matrices, for 14 MSCI indices from 11/08/2000 to 28/12/2010 (2610 days). Starting on 10/08/2000 ( $T = 1$ ), the last 4 years of data were used to estimate the 3 models, after which the estimates were used to produce rolling forecasts for the next 5 days. The model parameters were re-estimated taking into account new data every 5 days for a total of 522 re-estimations and 2610 out-of-sample forecasts. The performance statistics reported are  $\bar{W}_T$  representing terminal wealth of a portfolio with a starting value of 1, the mean ( $\hat{\mu}$ ), average volatility ( $\hat{\sigma}$ ), the annualized risk-return ( $RR = \sqrt{(252)} (\hat{\mu}/\hat{\sigma})$ ), the statistic and *p*-value of the Ledoit and Wolf (2008) test for the difference in the RR ratio between the IFACD( $\rho_t, \zeta_t$ ) and other models, the *p*-value of the MCS procedure of Hansen et al. (2011) using as loss function the negative of the portfolio returns and 10,000 bootstrap replications. The log(Relative Wealth) is the log of the ratio between the terminal wealth of the model (at time  $T$ ) and the terminal wealth obtained with the IFACD( $\rho_t, \zeta_t$ ) model. The CARA utility was optimized under 4 different risk aversion levels, from the mild ( $\eta = 1$ ) to very risk averse investor ( $\eta = 25$ ).

## References

- Barndorff-Nielsen, O. (1977). Exponentially decreasing distributions for the logarithm of particle size. *Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences*, 353:401–419.
- Barone Adesi, G., Gagliardini, P., and Urga, G. (2004). Testing asset pricing models with coskewness. *Journal of Business and Economic Statistics*, 22(4):474–485.
- Bauwens, L. and Laurent, S. (2005). A new class of multivariate skew densities, with application to generalized autoregressive conditional heteroscedasticity models. *Journal of Business and Economic Statistics*, 23(3):346–354.
- Beaulieu, M., Dufour, J., and Khalaf, L. (2007). Multivariate tests of mean-variance efficiency with possibly non-Gaussian errors. *Journal of Business and Economic Statistics*, 25(4):398–410.
- Beaulieu, M., Dufour, J., and Khalaf, L. (2009). Finite sample multivariate tests of asset pricing models with coskewness. *Computational Statistics & Data Analysis*, 53(6):2008–2021.
- Berkowitz, J. (2001). Testing density forecasts, with applications to risk management. *Journal of Business and Economic Statistics*, 19(4):465–474.
- Bläesild, P. (1981). The two-dimensional hyperbolic distribution and related distributions, with an application to Johanssen’s bean data. *Biometrika*, 68(1):251.
- Boswijk, H. P. and van der Weide, R. (2006). Wake me up before you GO-GARCH. Discussion Paper TI 2006-079/4, Tinbergen Institute.
- Boswijk, P. H. and van der Weide, R. (2011). Method of moments estimation of GO-GARCH models. *Journal of Econometrics*, 163(1):118–126.
- Broda, S. and Paoletta, M. (2009). CHICAGO: A fast and accurate method for portfolio risk calculation. *Journal of Financial Econometrics*, 7(4):412.
- Chabi-Yo, F., Ghysels, E., and Renault, E. (2008). On portfolio separation theorems with heterogeneous beliefs and attitudes towards risk. Working Paper/Document de travail 16, Bank of Canada.
- Chen, Y., Härdle, W., and Jeong, S. (2008). Nonparametric risk management with generalized hyperbolic distributions. *Journal of the American Statistical Association*, 103(483):910–923.
- Chen, Y., Härdle, W., and Spokoiny, V. (2007). Portfolio value at risk based on independent component analysis. *Journal of Computational and Applied Mathematics*, 205(1):594–607.
- Christoffersen, P. (1998). Evaluating interval forecasts. *International Economic Review*, 39(4):841–862.
- Christoffersen, P. and Pelletier, D. (2004). Backtesting value-at-risk: A duration-based approach. *Journal of Financial Econometrics*, 2(1):84–108.
- Comon, P. (1994). Independent component analysis, a new concept? *Signal Processing*, 36(3):287–314.
- Engle, R., Ng, V., and Rothschild, M. (1990). Asset pricing with a factor ARCH covariance structure: empirical estimates for Treasury bills. *Journal of Econometrics*, 45(1-2):213–237.
- Ferreira, J. and Steel, M. (2006). On describing multivariate skewed distributions: a directional approach. *Canadian Journal of Statistics*, 34(3):411–429.

- Hansen, B. (1994). Autoregressive conditional density estimation. *International Economic Review*, 35(3):705–730.
- Hansen, P., Lunde, A., and Nason, J. (2011). The model confidence set. *Econometrica*, 79(2):453–497.
- Harvey, C. and Siddique, A. (1999). Autoregressive conditional skewness. *Journal of Financial and Quantitative Analysis*, 34(4):465–487.
- Hyvärinen, A. and Oja, E. (2000). Independent component analysis: algorithms and applications. *Neural Networks*, 13(4-5):411–430.
- Jondeau, E. and Rockinger, M. (2003). Conditional volatility, skewness, and kurtosis: existence, persistence, and comovements. *Journal of Economic Dynamics and Control*, 27(10):1699–1737.
- Jondeau, E. and Rockinger, M. (2006). Optimal portfolio allocation under higher moments. *European Financial Management*, 12(1):29–55.
- Jondeau, E. and Rockinger, M. (2009). The impact of shocks on higher moments. *Journal of Financial Econometrics*, 7(2):77–105.
- Jurczenko, E. and Maillet, B. (2006). Theoretical foundations of asset allocation and pricing models with higher-order moments. In Jurczenko, E. and Maillet, B., editors, *Multi-moment Asset Allocation and Pricing Models*, pages 1–36. John Wiley & Sons.
- Kroner, K. and Ng, V. (1998). Modeling asymmetric comovements of asset returns. *Review of Financial Studies*, 11(4):817–844.
- Kupiec, P. (1995). Techniques for verifying the accuracy of risk measurement models. *Journal of Derivatives*, 3(2):73–84.
- Ledoit, O. and Wolf, M. (2008). Robust performance hypothesis testing with the sharpe ratio. *Journal of Empirical Finance*, 15(5):850–859.
- Markowitz, H. (1952). Portfolio selection. *Journal of Finance*, 7(1):77–91.
- Martellini, L. and Ziemann, V. (2010). Improved estimates of higher-order comoments and implications for portfolio selection. *Review of Financial Studies*, 23(4):1467–1502.
- Mencia, J. and Sentana, E. (2009). Multivariate location-scale mixtures of normals and mean-variance-skewness portfolio allocation. *Journal of Econometrics*, 153(2):105–121.
- Mencia, J. and Sentana, E. (2012). Distributional tests in multivariate dynamic models with normal and student innovations. *Review of Economics and Statistics*, 94(1):133–152.
- Rockinger, M. and Jondeau, E. (2002). Entropy densities with an application to autoregressive conditional skewness and kurtosis. *Journal of Econometrics*, 106(1):119–142.
- Ross, S. (1976). The arbitrage theory of capital asset pricing. *Journal of Economic Theory*, 13(3):341–360.
- Schmidt, R., Hrycej, T., and Stützle, E. (2006). Multivariate distribution models with generalized hyperbolic margins. *Computational Statistics & Data Analysis*, 50(8):2065–2096.
- Scott, R. and Horvath, P. (1980). On the direction of preference for moments of higher order than the variance. *Journal of Finance*, 35(4):915–919.

*The IFACD model*

- Sentana, E. (2009). The econometrics of mean-variance efficiency tests: a survey. *Econometrics Journal*, 12(3):C65–C101.
- van der Weide, R. (2002). GO-GARCH: a multivariate generalized orthogonal GARCH model. *Journal of Applied Econometrics*, 17(5):549–564.
- White, H. (1982). Maximum likelihood estimation of misspecified models. *Econometrica*, 50(1):1–25.
- Wilhelmsson, A. (2009). Value at risk with time varying variance, skewness and kurtosis—the NIG-ACD model. *Econometrics Journal*, 12(1):82–104.
- Zhang, K. and Chan, L. (2009). Efficient factor GARCH models and factor-DCC models. *Quantitative Finance*, 9(1):71–91.



## APPENDIX A: STANDARDIZED GH DENSITY

In order to model zero-mean, unit variance processes, the distribution, which must possess the scaling property, needs to be properly standardized. In the case of the GH distribution, because of the existence of location and scale invariant parameterizations and the possibility of expressing the mean and the variance in terms of one of those parameterizations, namely the  $(\zeta, \rho)$ , the task of standardizing the density can be broken down to one of estimating those 2 parameters, representing a combination of shape and skewness, followed by a series of transformation steps to translate the parameters into the  $(\alpha, \beta, \delta, \mu)$  parametrization for which standard formulae of the density function exist. The  $(\xi, \chi)$  parametrization, which is a simple transformation of the  $(\zeta, \rho)$ , could also be used in the first step and then transformed into the latter before proceeding further. In any case, moving between any of these parameterizations is a simple matter of applying the appropriate transformation. The steps to transforming from the  $(\zeta, \rho)$  to the  $(\alpha, \beta, \delta, \mu)$  parametrization, while at the same time standardizing for zero mean and unit variance are given henceforth. Let  $X$  be a random variable distributed as a  $GH(\zeta, \rho)$ , where

$$\zeta = \delta\sqrt{\alpha^2 - \beta^2}, \quad \rho = \frac{\beta}{\alpha}, \quad (29)$$

inverting (29) we can express  $\alpha$  and  $\beta$  in terms of  $\zeta$ ,  $\rho$  and  $\delta$

$$\alpha = \frac{\zeta}{\delta\sqrt{1 - \rho^2}}, \quad (30)$$

$$\beta = \alpha\rho. \quad (31)$$

For standardization we require that

$$\begin{aligned} E(X) &= \mu + \frac{\beta\delta}{\sqrt{\alpha^2 - \beta^2}} \frac{K_{\lambda+1}(\zeta)}{K_{\lambda}(\zeta)} = \mu + \frac{\beta\delta^2}{\zeta} \frac{K_{\lambda+1}(\zeta)}{K_{\lambda}(\zeta)} = 0 \\ Var(X) &= \delta^2 \left( \frac{K_{\lambda+1}(\zeta)}{\zeta K_{\lambda}(\zeta)} + \frac{\beta^2}{\alpha^2 - \beta^2} \left( \frac{K_{\lambda+2}(\zeta)}{K_{\lambda}(\zeta)} - \left( \frac{K_{\lambda+1}(\zeta)}{K_{\lambda}(\zeta)} \right)^2 \right) \right) = 1 \end{aligned} \quad (32)$$

it follows that

$$\mu = -\frac{\beta\delta^2}{\zeta} \frac{K_{\lambda+1}(\zeta)}{K_{\lambda}(\zeta)} \quad (33)$$

$$\delta = \left( \frac{K_{\lambda+1}(\zeta)}{\zeta K_{\lambda}(\zeta)} + \frac{\beta^2}{\alpha^2 - \beta^2} \left( \frac{K_{\lambda+2}(\zeta)}{K_{\lambda}(\zeta)} - \left( \frac{K_{\lambda+1}(\zeta)}{K_{\lambda}(\zeta)} \right)^2 \right) \right)^{-0.5} \quad (34)$$

Since we can express,  $\beta^2 / (\alpha^2 - \beta^2)$  as

$$\frac{\beta^2}{\alpha^2 - \beta^2} = \frac{\rho^2}{(1 - \rho^2)}, \quad (35)$$

then we can rewrite the formula for  $\delta$  in terms of the parameters  $\zeta$  and  $\rho$  as

$$\delta = \left( \frac{K_{\lambda+1}(\zeta)}{\zeta K_{\lambda}(\zeta)} + \frac{\rho^2}{(1 - \rho^2)} \left( \frac{K_{\lambda+2}(\zeta)}{K_{\lambda}(\zeta)} - \left( \frac{K_{\lambda+1}(\zeta)}{K_{\lambda}(\zeta)} \right)^2 \right) \right)^{-0.5} \quad (36)$$

Transforming into the  $(\alpha, \beta, \delta, \mu)$  parametrization proceeds by first substituting (36) into (30) and simplifying

$$\begin{aligned} \alpha &= \frac{\zeta \left( \frac{K_{\lambda+1}(\zeta)}{\zeta K_{\lambda}(\zeta)} + \frac{\rho^2 \left( \frac{K_{\lambda+2}(\zeta)}{K_{\lambda}(\zeta)} - \frac{(K_{\lambda+1}(\zeta))^2}{(K_{\lambda}(\zeta))^2} \right)}{(1 - \rho^2)} \right)^{0.5}}{\sqrt{(1 - \rho^2)}}, \\ &= \left( \frac{\zeta K_{\lambda+1}(\zeta)}{K_{\lambda}(\zeta)} \frac{1}{(1 - \rho^2)} \left( 1 + \frac{\zeta \rho^2 \left( \frac{K_{\lambda+2}(\zeta)}{K_{\lambda+1}(\zeta)} - \frac{K_{\lambda+1}(\zeta)}{K_{\lambda}(\zeta)} \right)}{(1 - \rho^2)} \right) \right)^{0.5}. \end{aligned} \quad (37)$$

Finally, the rest of the parameters are derived recursively from  $\alpha$  and the previous results

$$\beta = \alpha \rho, \quad (38)$$

$$\delta = \frac{\zeta}{\alpha \sqrt{1 - \rho^2}}, \quad (39)$$

$$\mu = \frac{-\beta \delta^2 K_{\lambda+1}(\zeta)}{\zeta K_{\lambda}(\zeta)}. \quad (40)$$

## APPENDIX B: THE GH CHARACTERISTIC FUNCTION

The moment generating function (MGF) of the GH Distribution is

$$\begin{aligned} M_{GH(\lambda, \alpha, \beta, \delta, \mu)}(u) &= e^{\mu u} M_{GIG}(\lambda, \delta \sqrt{\alpha^2 - \beta^2}) \left( \frac{u^2}{2} + \beta u \right), \\ &= e^{\mu u} \left( \frac{\alpha^2 - \beta^2}{\alpha^2 - (\beta + u)^2} \right)^{\lambda/2} \frac{K_{\lambda} \left( \delta \sqrt{\alpha^2 - (\beta + u)^2} \right)}{K_{\lambda} \left( \delta \sqrt{\alpha^2 - \beta^2} \right)} \end{aligned} \quad (41)$$

where  $M_{GIG}$  represents the moment generating function of the GIG which forms the mixing distribution in this variance-mean mixture subclass. The standardized skewness and kurtosis for the NIG

distribution have the following simplified expressions

$$\begin{aligned} K &= 3 + \frac{3(1 + 4\beta^2/\alpha^2)}{\delta\gamma} \\ S &= \frac{3\beta}{\alpha\sqrt{\delta\gamma}}. \end{aligned} \tag{42}$$

Powers of the MGF,  $M_{GH}(u)^p$ , only have the representation in (41) for  $p = 1$ , which means that GH distributions are not closed under convolution with the exception of the NIG, and only in the case when the shape and skew parameters are the same for all assets. When the distribution is not closed under convolution, numerical methods are required such as the inversion of the characteristic function by FFT.

Because the MGF is a holomorphic function for complex  $z$ , with  $|z| < \alpha - \beta$ , we can obtain the characteristic function of the GH distribution, using the following representation

$$\phi_{GH}(u) = M_{GHYP}(iu), \tag{43}$$

so that the characteristic function may be written as

$$\phi_{GH(\lambda, \alpha, \beta, \delta, \mu)}(u) = e^{\mu iu} \left( \frac{\alpha^2 - \beta^2}{\alpha^2 - (\beta + iu)^2} \right)^{\lambda/2} \frac{K_\lambda \left( \delta \sqrt{\alpha^2 - (\beta + iu)^2} \right)}{K_\lambda \left( \delta \sqrt{\alpha^2 - \beta^2} \right)}. \tag{44}$$

In order to find the portfolio density in the case of the IFACD model, the characteristic function required for the inversion of the NIG density is given below

$$\phi_{port}(u) = \exp \left\{ iu \sum_{j=1}^d \bar{\mu}_j + \sum_{j=1}^d \bar{\delta}_j \left( \sqrt{\bar{\alpha}_j^2 - \bar{\beta}_j^2} - \sqrt{\bar{\alpha}_j^2 - (\bar{\beta}_j + iu)^2} \right) \right\}, \tag{45}$$

where  $\bar{\alpha}_j$ ,  $\bar{\beta}_j$ ,  $\bar{\delta}_j$  and  $\bar{\mu}_j$  represent the parameters scaled as described in the main text of the paper. In the case of the GH characteristic function, this is a little more complicated as it involves the evaluation

of modified Bessel function of the third kind with complex arguments.<sup>3</sup> Taking logs and summing

$$\phi_{port}(u) = \exp \left\{ iu \sum_{j=1}^d \left( \bar{\mu}_j + \frac{\lambda_j}{2} \log (\bar{\alpha}_j^2 - \bar{\beta}_j^2) - \frac{\lambda_j}{2} \log (\bar{\alpha}_j^2 - (\bar{\beta}_j + iu)^2) + \right. \right. \\ \left. \left. \log \left( K_{\lambda_j} \left( \bar{\delta}_j \sqrt{\bar{\alpha}_j^2 - (\bar{\beta}_j + iu)^2} \right) \right) - \log \left( K_{\lambda_j} \left( \bar{\delta}_j \sqrt{\bar{\alpha}_j^2 - \bar{\beta}_j^2} \right) \right) \right) \right\} \quad (46)$$

which is more than 30 times slower to evaluate than the equivalent NIG function because of the Bessel function evaluations.

---

<sup>3</sup>Routines for this exist for example on netlib, see <http://www.netlib.org/amos/zbesk.f>